



GROUP 3 INSECTICIDE

RESTRICTED USE PESTICIDE
DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS.
For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF
AL, AR, DE, GA, IA, IL, IN, KS, KY, MA, MD, MN, MS, NJ, NC, NH, OH, PA, SC, TN, VA and WV



EPA Reg. No. 59639-35
FIFRA Section 2(ee) Recommendation
For use on Brown Marmorated Stink Bug

DIRECTIONS FOR USE

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. Always read and follow all label directions, restrictions and precautions when using any pesticide alone or in tank-mix combinations. The most restrictive labeling applies when using a tank mix.

RECOMMENDED FOR BROWN MARMORATED STINK BUG CONTROL

Pests	Crops	Danitol® 2.4 EC Spray Application Rates	Special Instructions
Brown Marmorated Stink Bug	Bushberries Cotton Cucurbits Head and Stem Brassica	10 2/3 to 16 fl oz/A (0.2 to 0.3 lb ai/A)	Apply <i>Danitol</i> 2.4 EC Spray as a foliar spray. Sufficient water volume should be used to provide complete and thorough coverage. Higher water volumes provide improved insect control. Insect control may be improved by the addition of a non-ionic surfactant to the spray solution according to manufacture's recommendation. Make applications beginning when first pest activity is noticed and repeat as needed. Under severe pest pressure, use the higher recommended rates. <i>Danitol</i> 2.4 EC Spray may be mixed with and/or alternated with commonly used insecticides to comply with local IPM programs. Continued on page 2
	Fruiting Vegetables Pea (Succulent)	10 2/3 fl oz/A (0.2 lb ai/A)	
	Grape Pome Fruit Stone Fruit Strawberry	10 2/3 to 21 1/3 fl oz/A (0.2 to 0.4 lb ai/A)	

Continued:

- Do not exceed 2 2/3 pt of *Danitol* 2.4 EC Spray (42 2/3 fl oz, 0.8 lb ai) per acre per season for all crops except bushberries and strawberry.
- Do not exceed more than two applications totaling 2 2/3 pt of *Danitol* 2.4 EC Spray (42 2/3 fl oz, 0.8 lb ai) per acre to the same planting in 12 consecutive months in strawberry.
- Do not exceed 2 pt of *Danitol* 2.4 EC Spray (32 fl oz, 0.6 lb ai) per acre per season in bushberries.
- Do not apply within twenty one (21) days of harvest for currant, cotton and grape.
- Do not apply within fourteen (14) days of harvest for pome fruit.
- Do not apply within seven (7) days of harvest for cucurbits, head and stem brassica and pea (succulent).
- Do not apply within three (3) days of harvest for bushberries, fruiting vegetables and stone fruit.
- Do not apply within two (2) days of harvest for strawberry.

APPLICATION INFORMATION AND PRECAUTIONS

- Always read and follow use directions, restrictions and precautions for all products. When tank mixing, the most restrictive labeling applies.
- Follow all applicable directions, restrictions and precautions on the EPA-registered labels.
- This labeling must be in the possession of the user at the time of pesticide application.
- It is a violation of Federal law to use these products in a manner inconsistent with their labeling.

Please contact Valent U.S.A. Corporation at 1-800-6-VALENT (682-5368) to determine if this product is registered in your state.

THIS RECOMMENDATION EXPIRES ON DECEMBER 31, 2012

Products That Work, From People Who Care[®] | www.valent.com | 800-6-VALENT (682-5368)

Registrant: Valent U.S.A. Corporation
P.O. Box 8025
Walnut Creek, CA 94596-8025

Danitol is a registered trademark of Sumitomo Chemical Company, Ltd.
Products That Work, From People Who Care is a registered trademark of Valent U.S.A. Corporation.

Product Bulletin



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Delegate[®] WG

EPA Reg. No. 62719-541

Special 2(ee) Recommendation[†]
(For Distribution and Use in All States Except Texas)

Suppression of Spotted Wing Drosophila on Various Crops

ATTENTION

[†]This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Delegate[®] WG insecticide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Delegate WG according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Delegate WG.

Directions for Use

Refer to product label for Delegate WG for General Use Precautions, Mixing Directions and Application Directions.

Delegate WG can be used as part of an integrated program to manage spotted wing drosophila. This use is limited to directed to ground application.

Apply Delegate WG to the following crops for suppression of spotted wing drosophila. Use a higher rate in the rate range for moderate to severe infestations and/or larger plant volume. Begin applications at first signs of adult activity. Due to the occurrence of multiple generations in a growing season, repeated applications may be required. Follow resistance management recommendations on the product label.

Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area. Refer to specific crop sections of the product label for additional information and restrictions.

Crop	Delegate WG (oz/acre)
bushberries (subgroup 13B)	3 - 6
caneberries (subgroup 13A)	
grape	3 - 5
pome fruits (crop group 11)	4.5 - 7
stone fruits (crop group 12)	

[®]Trademark of Dow AgroSciences LLC

R359-018

Issued: 08/18/10

Replaces R359-017

Torque™ Fungicide - Product Bulletin

EPA Reg. No. 69361-27-1001

FIFRA Section 2ee Recommendation - For Preventative and Curative Control of Fairy Ring caused by various Basidiomycete fungi

For distribution and use only within States where Torque Fungicide is currently registered (except NY and TX).

Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. The following recommendation is made as permitted under FIFRA Section 2(ee)(3) and has not been submitted to or approved by the US EPA.

Torque Fungicide may be used for the Preventative and Curative Control of Fairy Ring caused by various Basidiomycete fungi.

For Preventative treatment:

Apply Torque Fungicide before fairy ring symptoms appear in spring, starting when 2 inch soil temperatures consistently average 55-60°F (13-16°C), as a foliar spray at the rate of 0.6 fluid ounces per 1,000 square feet. Use sufficient water volume (66-132 GPA) and appropriate spray nozzles for thorough plant coverage. Water the application into the active root zone within 4 hours for best results. Do not tank-mix a wetting agent with preventive applications. Only use a wetting agent as required for the penetration of active hydrophobic soil conditions. User may make a second application of Torque in the same manner utilizing a 28-day interval. Torque may also be tank mixed or rotated with Affirm as allowed by label.

For Curative treatment:

Apply Torque Fungicide as a foliar spray at the rate of 0.6 fluid ounces per 1,000 square feet to areas where fairy ring symptoms are present. Use sufficient water volume (66-132 GPA) and appropriate spray nozzles for thorough plant coverage. Water the application into the active root zone within 4 hours for best results. Use a wetting agent as required for the penetration of active hydrophobic soil conditions. User may make a second application of Torque in the same manner utilizing a 28-day interval. Torque may also be tank mixed or rotated with Affirm as allowed by label.

Note: For First aid and other Precautionary statements, see label attached to the container. This recommendation is made as permitted under FIFRA Section 2(ee)(3) and has not been submitted to or approved by EPA. All applicable directions, restrictions, and precautions on the EPA registered label are to be followed. This bulletin must be in the possession of the user or applicator at the time of pesticide use.

Cleary Chemicals, LLC
178 Ridge Road, Suite A, Dayton, NJ 08810
(800) 524-1662
www.clearychemical.com

Torque™ is the Trademark of Cleary Chemicals, LLC.
©2010



Count on Cleary

syngenta

FIFRA

Date: September 21, 2010

Products: Concert® Fungicide
EPA Registration No. 100-1192

Use: For use on Gray Snow Mold (*Typhula incarnata* & *Typhula ishikariensis*) on turf

States: CO, CT, DE, IA, ID, IL, IN, MA, ME, MI, MN, MD, MT, NC, NE, ND, NH, NJ, OH, PA, RI, SD, UT, VA, VT, WA, WI, WV, WY

This recommendation is made under the authority of FIFRA Section 2(ee) and recommends an application of Concert Fungicide to turf at the recommended rates in the above states.

The user must refer to the federally approved labels for the above noted product and read and follow all directions for use, restrictions, and precautions.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

1. Use 8.5 fl. oz/1,000 sq. ft. of Concert Fungicide on turf. Apply at 14-28 day intervals.
2. Apply when conditions are favorable for disease development.

The user must refer to the federally approved labels for the above noted product and read and follow all directions for use, restrictions, and precautions.

The user should have this recommendation in its possession at the time of use.

Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist.

This recommendation for use of Concert Fungicide is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.

© 2010 Syngenta.

Syngenta Crop Protection, Inc.
Greensboro, NC 27419

Concert® and the Syngenta logo trademarks of a Syngenta group company.

Label code: CO1192096AC0910