



# Recommendation

FIFRA SECTION 2(ee) RECOMMENDATION FOR DISTRIBUTION AND USE IN THE STATES OF INDIANA, ILLINOIS, IOWA, KANSAS, KENTUCKY, MICHIGAN, MINNESOTA, MISSOURI, OHIO, PENNSYLVANIA, SOUTH DAKOTA, VIRGINIA, AND WISCONSIN

## TRICOR<sup>®</sup> DF 75% Dry Flowable Herbicide EPA Reg. No. 70506-103

### FOR FALL APPLICATION TO CORN AND SOYBEANS

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by USEPA.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read and follow all applicable instructions, restrictions, and precautions on the EPA-approved label. This label and the federally registered label must be in the possession of the user at the time of pesticide application.

WEEDS CONTROLLED	RATE: OUNCES TRICOR 75DF/A (Rate pounds ai/A)
Common Chickweed, Field Pennycress, Henbit, Marestalk, Winter Annual Mustard spp, Palmer Amaranth, Prickly Lettuce, Purple Deadnettle, Shepherd's Purse, Yellow Rocket, common dandelion seedlings	4- 12 (0.19 - 0.56)
<b>SPECIFIC DIRECTIONS</b>	
<p>Tricor™ 75% DF Herbicide may be used alone or in combination with other registered herbicides as a fall applied ground broadcast application for burndown and residual control. This fall application is made after the preceding crop has been harvested and at the first sign of germination of any of the listed winter annual broadleaf weeds.</p> <p>This application will reduce weed cover before spring planting. A fall application of TriCor 75DF is not intended to provide season long weed control. It is to be considered a part of a weed management program that will require additional application of a residual and/or post emergence herbicide for season-long control. This application will provide burndown and residual control of the listed germinating weeds. For optimum control, weeds must be less than 2 inches in height or diameter. For best results, make the application when the winter annual weeds begin germination.</p> <p>The length of residual control will increase with the application rate of TriCor 75DF. If emerged weeds are present and are greater than 2 inches in height or diameter, use 2,4-D or an appropriate alternative post emergence herbicide in a tank mixture with TriCor 75DF. To obtain maximum burndown of existing weeds of any size, use crop oil concentrate (COC) or an adjuvant in the tank mixture. Control of established common dandelion requires a tank mixture containing at least 1 pint/acre of a 4 pound/gallon of 2,4-D herbicide.</p> <p>Soybeans can be planted at any normal time the following spring. Corn can also be planted at any normal time the following spring after fall TriCor 75DF rates of 5 ½ ounces per acre or less. Corn can be planted at 4 or more months after fall Tricor 75DF at the rates on this document greater than 5.33 ounces per acre.</p>	

Rev. 11/14/12



**METRIBUZIN 75DF**  
EPA REG. NO.  
66222-106

**GLORY™**  
EPA REG. NO.  
66222-106

**GLORY™**  
★ HERBICIDE ★

### FIFRA §2(ee) RECOMMENDATION

For Fall or Early Spring Application in Corn or Soybeans and for Applications in Iowa, Illinois, Indiana, Kansas, Kentucky, Michigan, Minnesota, Missouri, Nebraska, Ohio, South Dakota and Wisconsin only

#### DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This recommendation must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered labels.
- This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by U.S. EPA.

Metribuzin 75DF and Glory can be used alone or in combination with other registered herbicides as part of a fall or spring applied (30 days before planting) herbicide program for burndown of existing vegetation prior to crop emergence in conservation tillage systems. Fall applications can be made anytime after harvest before ground freeze-up and up to 30 days prior to planting in the spring. Metribuzin 75DF or Glory may be tank mixed with "Roundup Brand" herbicides, Parazone 3SL, and other herbicides containing paraquat-dichloride for control of emerged weeds prior to field corn or soybean emergence. Metribuzin 75DF or Glory tank mixes with 2,4-DB, Fusion, Poast Plus, or Select may also be used in soybeans for control of emerged weeds prior to crop emergence. Metribuzin 75DF or Glory burndown tank mixes can be applied before planting or prior to crop emergence in the following areas:

**Field Corn:** Illinois, Indiana, Iowa, Kansas, Kentucky, Michigan, Minnesota, Missouri, Nebraska, Ohio, South Dakota, and Wisconsin.

**Soybeans:** All areas for all products except Fusion tank mixes. See Fusion section on the production label for recommended states.

**Application:** Spring applications of Metribuzin 75DF and Glory may be applied up to 30 days prior to planting or preemergence. Apply only by ground equipment when Metribuzin 75DF is used for burndown of existing vegetation in conservation tillage systems. Metribuzin 75DF and Glory tank mix partner burndown rates are listed in the following three tables.

METRIBUZIN 75 DF AND GLORY BURNDOWN RATES Field Corn and Soybeans		
Crops	Application Timing	Metribuzin 75DF Rate (oz./A)
Field Corn: Iowa, Kansas, Missouri, Nebraska, South Dakota	Preplant (0 to 30 days)	2 to 5½
	Preemergence	
Field corn: Illinois, Indiana, Kentucky, Michigan, Minnesota, Ohio, Wisconsin	Preplant (10 to 30 days)	2 to 5½
	Preplant (0 to 9 days)	2 to 4
	Preemergence	
Soybeans	Preplant (0 to 30 days)	2 to 5½
	Preemergence	

**SPECIAL PRECAUTIONS:** Do not apply these treatments after crop emergence. Observe all precautions and limitations on the labeling of all products used in tank mixtures. Refer to the GENERAL INFORMATION section of this label for additional information, precautions, and limitations.

**Field Corn:**

1. Do not apply on coarse-textured soils with less than 1.5% organic matter.
2. Do not apply more than 4 oz. of Metribuzin 75DF or Glory per acre on soils with less than 2% organic matter.
3. Do not apply on soils having pH 7.0 or greater.
4. Do not apply more than 5½ oz. Metribuzin 75DF or Glory (0.25 pound active ingredient) per acre per growing season.
5. Corn seed should be planted a minimum of 1½ inches deep.
6. Metribuzin 75DF or Glory may only be used in hybrid seed corn production fields if both inbred parents are known to be tolerant to Metribuzin 75DF.

**Soybeans:**

1. Apply only 2,4-D low volatile ester formulations which are registered and recommended for preplant or burndown use in soybeans.
2. Do not apply tank mixture containing 2,4-D LVE if wind is blowing toward desired susceptible plants (i.e. cotton, tobacco, tomato, etc.) or when wind speeds exceed 6 miles per hour.

**Feeding Restrictions:** Corn treated with Metribuzin 75DF or Glory may be harvested for silage or grain 60 days after treatment. Soybean vines or hay treated with Metribuzin 75DF or Glory may be grazed or fed to livestock 40 days after application. Do not feed hay, forage, fodder, or graze 2,4-D-, Select-, or Fusion-treated vegetation. Follow the most restrictive preharvest interval of all products used in a tank mixture.

Read the "LIMITATION OF WARRANTY AND LIABILITY" in the label booklet for Metribuzin 75DF and Glory before using.

10/01/12  
E 110504/092605/051309/N 060508

