



FIFRA

Section 2(ee) Recommendation

Date: February 22, 2016

Product: Tenacity® (EPA Reg. No. 100-1267)
+
Turflon® Ester Ultra (EPA Reg. No. 62719-566)

Use: Addition of a Tank Mix and Addition of Pests for Control of Broadleaf Species in Certain Turfgrass Varieties

State: AR, CA, CO, CT, DC, DE, FL, GA, HI, IA, ID, IL, IN, KS, KY, LA, MA, ME, MI, MN, MO, MS, MT, NC, ND, NE, NH, NJ, NM, NV, OH, OK, PA, RI, SC, SD, TN, UT, VA, VT, WA, WI, WV, WY

Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- The user must refer to the federally approved labels for the above noted product and read and follow all directions for use, restrictions, and precautions.
- The user should have this recommendation in its possession at the time of use.
- Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist.
- This recommendation for use of this product is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.

This 2(ee) Recommendation applies to the following turf species only: tall fescue, Kentucky Bluegrass, and perennial ryegrass. **Additional pests controlled:** black medic (*Medicago lupulina*) and common bermudagrass.

For postemergence control of listed broadleaf species, apply Tenacity at 5 fl oz/acre plus Turflon Ester Ultra at 16 fl oz/acre. Add a non-ionic surfactant at 0.25 to 0.5% v/v. Make 2 applications on a 4-6 week interval.

For common bermudagrass control, apply Tenacity at 5 fl oz/acre plus Turflon Ester Ultra at 16 fl oz/acre. Add a non-ionic surfactant at 0.25 to 0.5% v/v. Make 3 applications on a 21 day interval when common bermudagrass is actively growing. Do not overseed for 21 days after the last application.

Follow all directions for use, precautions, and restrictions on the federally approved Tenacity and Turflon Ester Ultra labels.

Tenacity® and the Syngenta logo are trademarks of a Syngenta Group Company.
Turflon® trademark of Dow AgroSciences LLC.

©2016 Syngenta

2(ee) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Expiration Date: February 22, 2020

Label Code: AR1267045AC0216

2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



DUPONT™ HARMONY® SG herbicide (with TotalSol® soluble granules)

EPA Reg. No. 352-633

DUPONT™ EXPRESS® herbicide (with TotalSol® soluble granules)

EPA Reg. No. 352-632

DUPONT™ ALLY® XP Herbicide

EPA Reg. No. 352-435

DUPONT™ DICAMBA XP Herbicide

EPA Reg. No. 7969-140-352

OR

DUPONT™ BL1 Herbicide

EPA Reg. No. 42750-271-352

FOR POSTEMERGENCE WEED CONTROL OF BROADLEAF WEEDS IN WHEAT, WINTER BARLEY, TRITICALE and FALLOW

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

HARMONY® SG herbicide (with TotalSol® soluble granules) may be tank mixed at 0.272 ounces per acre with EXPRESS® herbicide (with TotalSol® soluble granules) at 0.137 ounces per acre plus ALLY® XP herbicide at 0.091 ounces per acre and 2.88 ounces per acre of Dicamba XP herbicide or DuPont™ BL1 herbicide for the control of broadleaf weeds when applied postemergence to winter wheat and winter barley in the fall or spring after the 2-leaf stage but prior to the jointing stage, when the majority of weeds have emerged and are actively growing. Spring applications to spring wheat must be made after the 2-leaf stage before the wheat exceeds the 6-leaf stage, when the majority of weeds have emerged and are actively growing. Fall applications to triticale should be applied after the 2-leaf stage but prior to jointing, while spring seeded triticale applications must be made after the 2-leaf stage before the triticale reaches the 6-leaf stage. Application of this combination should not be made to small grains underseeded to legumes as severe legume injury or death will occur. Fallow applications should be made in the spring, summer, or fall when the majority of weeds have emerged and are actively growing. Refer to all labels for an entire lists of weeds controlled.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or DuPont 2(ee) instructions.

Read and follow all applicable directions, restrictions and precautions on the EPA-registered label for each product.

© 2015 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A

Unless indicated; trademarks with ®; TM or SM are trademarks of DuPont or its affiliates. ©2015 DuPont

IMPORTANT PRECAUTIONS:

Refer to the HARMONY® SG herbicide (with TotalSol® soluble granules), EXPRESS® herbicide (with TotalSol® soluble granules), ALLY® XP herbicide, Dicamba XP herbicide, DuPont™ BL1 herbicide and any tank mix partner labels for information on weed species controlled, additional use directions, restrictions, rotational crop intervals and precautions. The most restrictive provision on any label shall apply.

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2015 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A

Unless indicated; trademarks with ®; TM or SM are trademarks of DuPont or its affiliates. ©2015 DuPont

2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



DUPONT™ EXPRESS® herbicide (with TotalSol® soluble granules)

EPA Reg. No. 352-632

DUPONT™ ALLY® XP herbicide

EPA Reg. No. 352-435

PLUS DUPONT™ GR1™

EPA REG. NO. 352-884

FOR POSTEMERGENCE WEED CONTROL OF BROADLEAF AND GRASSY WEEDS IN WINTER WHEAT

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

EXPRESS® herbicide (with TotalSol® soluble granules) may be tank mixed at 0.27 ounces per acre with ALLY® XP herbicide at 0.03 ounces per acre plus GR1™ at 2.0 ounces per acre for control of broadleaf and grassy weeds when applied postemergence to winter wheat in the 3-leaf to jointing stage of growth, for the control of annual broadleaf and grassy weeds. Refer to all labels for an entire lists of weeds controlled.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or DuPont 2(ee) instructions.

Read and follow all applicable directions, restrictions and precautions on the EPA-registered label for each product.

IMPORTANT PRECAUTIONS:

Refer to the EXPRESS® herbicide (with TotalSol® soluble granules), ALLY® XP herbicide, GR1™ and any tank mix partner labels for information on weed species controlled, additional use directions, restrictions, rotational crop intervals and precautions. The most restrictive provision on any label shall apply.

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application. Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2015 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A

R-1671 111615 11-20-15 C12

Page 1 of 1

Expires 12/31/18

2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



DUPONT™ EXPRESS® herbicide (with TotalSol® soluble granules)

EPA Reg. No. 352-632

DUPONT™ ALLY® XP Herbicide

EPA Reg. No. 352-435

FOR POSTEMERGENCE WEED CONTROL OF BROADLEAF WEEDS IN WHEAT, BARLEY, TRITICALE and FALLOW

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

EXPRESS® herbicide (with TotalSol® soluble granules) may be tank mixed at 0.37 ounces per acre plus ALLY® XP herbicide at 0.03 ounces per acre for the control of broadleaf weeds when applied postemergence to winter wheat and triticale in the fall after the 2-leaf stage but before the flag leaf is visible, when the majority of weeds have emerged and are actively growing. Spring applications to winter wheat, spring wheat, barley, and triticale must be made after the 2-leaf stage but before the flag leaf is visible, when the majority of weeds have emerged and are actively growing. Application of this combination should not be made to small grains underseeded to legumes as severe legume injury or dead will occur. Fallow applications should be made in the spring, summer, or fall when the majority of weeds have emerged and are actively growing. Refer to all labels for an entire lists of weeds controlled.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or DuPont 2(ee) instructions.

Read and follow all applicable directions, restrictions and precautions on the EPA-registered label for each product.

IMPORTANT PRECAUTIONS:

Refer to the EXPRESS® herbicide (with TotalSol® soluble granules), ALLY® XP herbicide, and any tank mix partner labels for information on weed species controlled, additional use directions, restrictions, rotational crop intervals and precautions. The most restrictive provision on any label shall apply.

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2015 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A

TECHNICAL INFORMATION BULLETIN

RANCONA[®] 3.8 FS
Seed Treatment Fungicide

EPA Reg. No. 400-544

FIFRA Section 2(ee) Recommendation

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF
WASHINGTON, OREGON, MONTANA, IDAHO, MINNESOTA, WISCONSIN
AND ILLINOIS**

This recommendation must be in the possession of the user at the time of application.

**Use on Beans and Peas (Succulent or Dried) as a seed treatment
to control *Ascochyta* seedling blight**

DIRECTION FOR USE

Use information in accordance with Section 2(ee) of FIFRA.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. **All applicable directions, restrictions, and precautions on the EPA-registered label for Rancona 3.8 FS are to be followed.** This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the cautionary statements plus all other information appearing on the Rancona 3.8 FS product label.

BEANS AND PEAS

Target Pest	Application Rate (fl oz of product per 100 lbs. of seed)	Application Instructions
Seedling blight (seed borne <i>Ascochyta</i> spp.)	0.085	Treat seed at the application rate for control of seed-borne <i>Ascochyta</i> blight.

MacDermid Agricultural Solutions, Inc.
245 Freight St.
Waterbury, CT 06702-1818

Rancona[®] is a registered trademark of MacDermid Agricultural Solutions, Inc..

Issue Date: February 3, 2015/6

TECHNICAL INFORMATION BULLETIN

RANCONA[®] CTS
Seed Protectant

EPA Reg. No. 400-592

FIFRA Section 2(ee) Recommendation

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF
WASHINGTON, OREGON, MONTANA, IDAHO, MINNESOTA, WISCONSIN
AND ILLINOIS**

This recommendation must be in the possession of the user at the time of application.

**Use on dried shelled and succulent / edible podded Beans and
Peas as a seed treatment to control *Ascochyta* seedling blight**

DIRECTION FOR USE

Use information in accordance with Section 2(ee) of FIFRA.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. All applicable directions, restrictions, and precautions on the EPA-registered label for Rancona CTS are to be followed. This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the cautionary statements plus all other information appearing on the Rancona CTS product label.

BEANS AND PEAS

Target Pest	Application Rate (fl oz product per 100 lbs. of seed)	Application Instructions
Seedling blight (seed borne <i>Ascochyta</i> spp.)	1.53	Treat seed at the application rate for control of seed-borne <i>Ascochyta</i> blight.

MacDermid Agricultural Solutions, Inc.
245 Freight St.
Waterbury, CT 06702-1818

Rancona ® is a registered trademark of MacDermid Agricultural Solutions, Inc.

Issue Date: February 2, 2015

TECHNICAL INFORMATION BULLETIN

RANCONA[®] Summit
Seed Protectant

EPA Reg. No. 400-570

FIFRA Section 2(ee) Recommendation

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF
WASHINGTON, OREGON, MONTANA, IDAHO, MINNESOTA, WISCONSIN
AND ILLINOIS**

This recommendation must be in the possession of the user at the time of application.

**Use on dried shelled and succulent / edible podded Beans and
Peas as a seed treatment to control *Ascochyta* seedling blight**

DIRECTION FOR USE

Use information in accordance with Section 2(ee) of FIFRA.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. **All applicable directions, restrictions, and precautions on the EPA-registered label for Rancona Summit are to be followed.** This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the cautionary statements plus all other information appearing on the Rancona Summit product label.

BEANS AND PEAS

Target Pest	Application Rate (fl oz of product per 100 lbs. of seed)	Application Instructions
Seedling blight (seed borne <i>Ascochyta</i> spp.)	4.0	Treat seed at the application rate for control of seed-borne <i>Ascochyta</i> blight.

MacDermid Agricultural Solutions, Inc.
245 Freight St.
Waterbury, CT 06702-1818

Rancona ® is a registered trademark of MacDermid Agricultural Solutions, Inc.

Issue Date: February 3, 2015/6

TECHNICAL INFORMATION BULLETIN

RANCONA[®] V 100 Pro FS
Seed Protectant

EPA Reg. No. 400-596

FIFRA Section 2(ee) Recommendation

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF
WASHINGTON, OREGON, MONTANA, IDAHO, MINNESOTA, WISCONSIN
AND ILLINOIS**

This recommendation must be in the possession of the user at the time of application.

**Use on dried shelled Beans as a seed treatment to control
Ascochyta seedling blight**

DIRECTION FOR USE

Use information in accordance with Section 2(ee) of FIFRA.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. All applicable directions, restrictions, and precautions on the EPA-registered label for Rancona V 100 Pro FS are to be followed. This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the cautionary statements plus all other information appearing on the Rancona V 100 Pro FS product label.

BEANS

Target Pest	Application Rate (fl oz product per 100 lbs. of seed)	Application Instructions
Seedling blight (seed borne <i>Ascochyta</i> spp.)	1.5	Treat seed at the application rate for control of seed-borne <i>Ascochyta</i> blight.

MacDermid Agricultural Solutions, Inc.
245 Freight St.
Waterbury, CT 06702-1818

Rancona[®] is a registered trademark of MacDermid Agricultural Solutions, Inc.

Issue Date: February 3, 2015/10

FIFRA Section 2(ee) Recommendation



GROUP	14	HERBICIDE
-------	----	-----------

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF
AL, AR, DE, GA, IA, IL, IN, KS, KY, LA, MD, MI, MN, MO, NC,
ND, NE, NJ, OH, OK, SC, SD, TN, VA and WI



EPA Reg. No. 59639-99

**VALOR[®] SX HERBICIDE PLUS BOUNDARY[®], ALACHLOR,
S-METOLACHLOR, DIMETHENAMID-P OR FLUFENACET HERBICIDES FOR EARLY
PRE-PLANT APPLICATIONS AT A MINIMUM OF 14 DAYS PRIOR TO PLANTING FOR
RESIDUAL CONTROL OF CERTAIN WEEDS IN NO-TILLAGE AND REDUCED-TILLAGE
SOYBEANS**

OR

**VALOR SX HERBICIDE APPLIED AT A MINIMUM OF 14 DAYS PRIOR TO PLANTING
FOLLOWED BY BOUNDARY, ALACHLOR, S-METOLACHLOR, DIMETHENAMID-P OR
FLUFENACET HERBICIDES APPLIED AT PLANTING FOR RESIDUAL CONTROL OF
CERTAIN WEEDS IN NO-TILLAGE AND REDUCED-TILLAGE SOYBEANS**

THIS FIFRA SECTION 2(E) EXPIRES DECEMBER 31, 2021

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This 2(ee) recommendation contains directions for use which do not appear on the package label and must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions and precautions on the EPA registered label when using *Valor SX* Herbicide. Always read and follow all label directions when using any pesticide alone or in tank mix combinations. The most restrictive labeling applies when using a tank mix.

- Do not graze animals on green forage or stubble. Do not utilize hay or straw for animal feed or bedding.

WEEDS CONTROLLED	ORGANIC MATTER	SOIL TYPE	RATES
Lambsquarters, Common Nightshades Black Eastern Black Pigweeds Redroot Palmer Amaranth Smooth Spiny Amaranth Tumble Sida, Prickly Velvetleaf ¹	Up to 5%	All Soil Types	<i>Valor</i> SX Herbicide at 1.5 to 3.0 oz/A plus Boundary 7.8 EC at 0.5 to 2 pt/A or Boundary 6.5 EC at 0.6 to 2.4 pt/A or alachlor at 1 to 2 lb ai/A or s-metolachlor at 0.5 to 2 lb ai/A or dimethenamid-p at 0.25 to 0.6 lb ai/A or flufenacet at 0.15 to 0.45 lb ai/A
<p>APPLICATION METHOD: These early preplant applications must be surface applied. Incorporation of these tank mixes will result in reduced control of the weeds listed above.</p> <p>NOTE: Refer to <i>Valor</i> SX Herbicide, Boundary 7.8 EC, Boundary 6.5 EC, alachlor, s-metolachlor, dimethenamid-p or flufenacet labels for other weeds controlled.</p>			

¹A postemergence herbicide, such as *Resource*[®] or glyphosate plus *Resource* (Roundup Ready[®] soybeans only) may be needed to adequately control velvetleaf.

Please contact Valent U.S.A. Corporation at 800-6-VALENT (682-5368) to determine if this product is registered in your state.

©2016 Valent U.S.A. Corporation

Resource and *Valor* are registered trademarks of Valent U.S.A. Corporation.
Boundary is a registered trademark of Syngenta, EPA Reg. No. 100-1162 264-771
Roundup Ready is a registered trademark of Monsanto Company.

Manufactured for:
Valent U.S.A. Corporation
P.O. Box 8025
Walnut Creek, CA 94596-8025

Made in U.S.A.

FIFRA Section 2(ee) Recommendation



GROUP	14	HERBICIDE
-------	----	-----------

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF
AL, AR, DE, GA, IA, IL, IN, KS, KY, LA, MD, MI, MO, NC, NE,
NJ, OH, OK, SC, TN, VA and WI



EPA Reg. No. 59639-117

**VALOR[®] XLT SOYBEAN HERBICIDE PLUS BOUNDARY[®], ALACHLOR,
S-METOLACHLOR, DIMETHENAMID-P OR FLUFENACET HERBICIDES FOR EARLY
PRE-PLANT APPLICATIONS AT A MINIMUM OF 14 DAYS PRIOR TO PLANTING FOR
RESIDUAL CONTROL OF CERTAIN WEEDS IN NO-TILLAGE AND REDUCED-TILLAGE
SOYBEANS**

OR

**VALOR XLT SOYBEAN HERBICIDE APPLIED AT A MINIMUM OF 14 DAYS PRIOR TO
PLANTING FOLLOWED BY BOUNDARY, ALACHLOR, S-METOLACHLOR,
DIMETHENAMID-P OR FLUFENACET HERBICIDES APPLIED AT PLANTING FOR
RESIDUAL CONTROL OF CERTAIN WEEDS IN NO-TILLAGE AND REDUCED-TILLAGE
SOYBEANS**

THIS FIFRA SECTION 2(EE) EXPIRES DECEMBER 31, 2021

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This 2(ee) recommendation contains directions for use which do not appear on the package label and must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions and precautions on the EPA registered label when using *Valor XLT Soybean Herbicide*. Always read and follow all label directions when using any pesticide alone or in tank mix combinations. The most restrictive labeling applies when using a tank mix.

- In the states of Delaware, Maryland, New Jersey and Virginia do not exceed 3.0 oz/A of *Valor XLT Soybean Herbicide*.
- Do not graze animals on green forage or stubble. Do not utilize hay or straw for animal feed or bedding.

WEEDS CONTROLLED	ORGANIC MATTER	SOIL TYPE	RATES
Lambsquarters, Common Nightshades Black Eastern Black Pigweeds Redroot Palmer Amaranth Smooth Spiny Amaranth Tumble Sida, Prickly Velvetleaf ¹	Up to 5%	All Soil Types	<i>Valor</i> XLT Soybean Herbicide at 3 to 5 oz/A plus Boundary 7.8 EC at 0.5 to 2 pt/A or Boundary 6.5 EC at 0.6 to 2.4 pt/A or alachlor at 1 to 2 lb ai/A or s-metolachlor at 0.5 to 2 lb ai/A or dimethenamid-p at 0.25 to 0.6 lb ai/A or flufenacet at 0.15 to 0.45 lb ai/A

APPLICATION METHOD: These early preplant applications must be surface applied. Incorporation of these tank mixes will result in reduced control of the weeds listed above.

NOTE: Refer to *Valor* XLT Soybean Herbicide, Boundary 7.8 EC, Boundary 6.5 EC, alachlor, s-metolachlor, dimethenamid-p or flufenacet labels for other weeds controlled.

¹A postemergence herbicide, such as *Resource*[®] or glyphosate plus *Resource* (Roundup Ready[®] soybeans only) may be needed to adequately control velvetleaf.

Please contact Valent U.S.A. Corporation at 800-6-VALENT (682-5368) to determine if this product is registered in your state.

©2016 Valent U.S.A. Corporation

Resource and *Valor* are registered trademarks of Valent U.S.A. Corporation.
Boundary is a registered trademark of Syngenta, EPA Reg. No. 100-1162 264-771
Roundup Ready is a registered trademark of Monsanto Company.

Manufactured for:
Valent U.S.A. Corporation
P.O. Box 8025
Walnut Creek, CA 94596-8025

Made in U.S.A.

RESTRICTED USE PESTICIDE

May injure (phytotoxic) susceptible non-target plants.

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Commercial and certified applicators must ensure that all persons involved in these activities are informed of the precautionary statements.



Bayer CropScience

Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

CORVUS® Herbicide

EPA Reg. No. 264-1066

CORVUS® Herbicide + Atrazine + Glyphosate tank for Early Postemergent use in corn to control of broadleaf and glyphosate-resistant weeds.

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF ARKANSAS, ILLINOIS, INDIANA, IOWA, KANSAS, KENTUCKY, MICHIGAN, MISSOURI, OHIO, SOUTH DAKOTA AND TENNESSEE

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. This product bulletin should be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling.

CORVUS® Herbicide can be applied to corn in tank mixture with atrazine and glyphosate from spiking through the 2-leaf collar growth stage.

Product	Rate	Crop	Weeds
CORVUS® Herbicide + atrazine + glyphosate	3.33 to 4.5 fl oz/acre ¹ + 0.5 lb ai/acre + 8-12 fl oz/acre of glyphosate	Apply tank mixture to field corn from spike up to 2-leaf collar growth stage	For enhanced control of broadleaf and glyphosate-resistant weeds

¹Refer to the Specific Use Directions table of the CORVUS® Herbicide label for use rates by soil texture.

Notes and Restrictions:

- For use only on glyphosate-tolerant corn.
- Early season crop response may be observed under certain environmental conditions.
- Do not use COC or MSO with CORVUS® Herbicide applied to emerged field corn.
- Tank-mixtures with other herbicides or adjuvants are not recommended for early postemergence applications of CORVUS® Herbicide.

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937)

RESTRICTED USE PESTICIDE

May injure (phytotoxic) susceptible non-target plants.

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Commercial and certified applicators must ensure that all persons involved in these activities are informed of the precautionary statements.



Bayer CropScience

Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

CORVUS® Herbicide

EPA Reg. No. 264-1066

CORVUS® Herbicide + Atrazine + Dicamba tank mixture for Early Postemergent use in corn to control of broadleaf and glyphosate-resistant weeds.

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF ARKANSAS, ILLINOIS, INDIANA, IOWA, KANSAS, KENTUCKY, MICHIGAN, MISSOURI, OHIO, SOUTH DAKOTA AND TENNESSEE

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. This product bulletin should be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling.

CORVUS® Herbicide can be applied to corn in tank mixture with atrazine and dicamba from spiking through the 2-leaf collar growth stage.

Product	Rate	Crop	Weeds
CORVUS® Herbicide	3.33 to 4.5 fl oz/acre ¹	Apply tank mixture to field corn from spike up to 2-leaf collar growth stage	For enhanced control of broadleaf and glyphosate-resistant weeds
+	+		
atrazine	0.5 lb ai/acre		
+	+		
dicamba	8 fl oz/acre		

¹Refer to the Specific Use Directions table of the CORVUS® Herbicide label for use rates by soil texture.

Notes and Restrictions:

- Early season crop response may be observed under certain environmental conditions.
- Do not use COC or MSO with CORVUS® Herbicide applied to emerged field corn.
- Tank-mixtures with other herbicides or adjuvants are not recommended for early postemergence applications of CORVUS® Herbicide.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937)



FIFRA

Section 2(ee) Recommendation

Date: January 7, 2016

Product: Orondis® Gold 200 (EPA Reg. No. 100-1571)
+
Ridomil® Gold SL (EPA Reg. No. 100-1202)

Use: **Addition of a Tank Mix of Orondis Gold 200 with Ridomil Gold SL**

State: AL, AR, AZ, CO, DE, FL, GA, IA, ID, IL, IN, KS, LA, MA, MD, ME, MI, MN, MO, MT, NC, ND, NE, NH, NJ, NM, NV, OH, OK, PA, SC, SD, TN, UT, VA, VT, WA, WI, WV, WY

Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- The user must refer to the federally approved labels for the above noted product and read and follow all directions for use, restrictions, and precautions.
- The user should have this recommendation in its possession at the time of use.
- Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist.
- This recommendation for use of this product is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.

For control of diseases on Cucurbits, Fruiting Vegetables (including Tomato), Leafy Greens/Leafy Vegetables (except Brassica) tank mix Orondis Gold 200 at 4.8 fl oz/A with Ridomil Gold SL at 0.5 pt/A (8 fl oz/A).

Shake Orondis Gold 200 container well before using. Follow all directions for use, restrictions and precautions given on the Orondis Gold 200 and Ridomil Gold SL labels.

Orondis® and the Syngenta logo are trademarks of a Syngenta Group Company.

©2016 Syngenta

2(ee) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Expiration Date: January 7, 2020

Label Code: AL1571127AC0116



Bayer CropScience

Bayer CropScience LP
 P.O. Box 12014
 2 T.W. Alexander Drive
 Research Triangle Park, North Carolina 27709
 1-866-99BAYER (1-866-992-2937)

SIVANTO™ prime

EPA Reg. No. 264-1141

For suppression of Woolly Apple Aphid in Pome Fruit.

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF ALABAMA, ALASKA, ARIZONA, ARKANSAS, CALIFORNIA, COLORADO, DELAWARE, DISTRICT OF COLUMBIA, FLORIDA, GEORGIA, IDAHO, ILLINOIS, INDIANA, IOWA, KANSAS, KENTUCKY, LOUISIANA, MAINE, MASSACHUSETTS, MICHIGAN, MISSISSIPPI, MISSOURI, MONTANA, NEBRASKA, NEW JERSEY, NORTH CAROLINA, NORTH DAKOTA, OHIO, OKLAHOMA, PENNSYLVANIA, RHODE ISLAND, SOUTH CAROLINA, SOUTH DAKOTA, TENNESSEE, UTAH, VERMONT, VIRGINIA, WASHINGTON, WEST VIRGINIA, WISCONSIN AND WYOMING

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. This product bulletin should be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

POME FRUIT – FOLIAR

Crops of Crop Group 11-10 Including: Apple, Azarole, Crabapples (Chinese apple, Chinese crab apple, Chinese flowering apple, Crab apple, Cutleaf crab apple, Florentine crab apple, Hall crab apple, Iowa crab apple, Japanese crab apple, Kai do crab apple, Manchurian crab apple, Paradise apple, Sargent's crab apple, Siberian crab apple, Soulard crab apple, Southern crab apple, Sweet crab apple, Tea crab apple, Toringa crab apple, Western Crabapple, Yunnan crab apple, and varieties and/or hybrids of these), Loquat, Mayhaw, Medlar, Pear, Asian pear, Quince, Chinese quince, Japanese quince, Tejocote, and cultivars, varieties and/or hybrids of these.

Pest Suppressed	Product Rate (fl oz/A)
Woolly apple aphid	12.0 – 14.0

Foliar Application Restrictions:

Pre-Harvest Interval (PHI): **14 day**
 Minimum interval between applications: **10 days**
 Minimum application volumes: **25 gallons/Acre (Ground); 10 gallons/Acre (Aerial)**
 Maximum SIVANTO prime allowed per year: **28.0 fluid ounces/Acre (0.365 lb AI/Acre).**

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

*As with any crop-protection product, always read and follow label directions.
 For additional information call toll-free 1-866-99BAYER (1-866-992-2937)*

Sivanto is a trademark of Bayer.



Bayer CropScience

Bayer CropScience LP
 P.O. Box 12014
 2 T.W. Alexander Drive
 Research Triangle Park, North Carolina 27709
 1-866-99BAYER (1-866-992-2937)

SIVANTO™ prime

EPA Reg. No. 264-1141

For suppression of Blueberry Thrips in Blueberry.

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF ALABAMA, ALASKA, ARIZONA, CALIFORNIA, COLORADO, DELAWARE, DISCTRICT OF COLUMBIA, FLORIDA, GEORGIA, IDAHO, ILLINOIS, INDIANA, IOWA, KANSAS, KENTUCKY, LOUISIANA, MAINE, MASSACHUSETTS, MICHIGAN, MISSISSIPPI, MISSOURI, MONTANA, NEBRASKA, NEW JERSEY, NORTH CAROLINA, NORTH DAKOTA, OHIO, OKLAHOMA, PENNSYLVANIA, RHODE ISLAND, SOUTH CAROLINA, SOUTH DAKOTA, TENNESSEE, UTAH, VERMONT, VIRGINIA, WASHINGTON, WEST VIRGINIA, WISCONSIN AND WYOMING

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. This product bulletin should be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BLUEBERRY	
Pest Suppressed	Product Rate (fl oz/A)
Blueberry thrips - <i>Scirtothrips citri</i>	10.5 – 14.0
Foliar Application Restrictions: Pre-Harvest Interval (PHI): 3 days Minimum interval between applications: 7 days Minimum application volumes: 25 gallons/Acre (Ground); 2 gallons/Acre (Aerial) Maximum SIVANTO prime allowed per year: 28.0 fluid ounces/Acre (0.365 lb AI/Acre).	

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937)



Bayer CropScience

Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

SIVANTO™ prime
EPA Reg. No. 264-1141

Reduced rates for foliar applications in Sorghum in control of Sugarcane Aphid.

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF ALABAMA, ARKANSAS, DELAWARE, FLORIDA, GEORGIA, ILLINOIS, KANSAS, KENTUCKY, LOUISIANA, MARYLAND, MISSISSIPPI, MISSOURI, NEBRASKA, NORTH CAROLINA, OKLAHOMA, SOUTH CAROLINA, TENNESSEE, AND VIRGINIA

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. This product bulletin should be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SORGHUM – FOLIAR

Pest Controlled	Product Rate (fl oz/A)
Sugarcane Aphid	4.0 – 7.0
Foliar Application Restrictions: Pre-Harvest Interval (PHI): 7 days - forage; 21 days - dried grain, stover or straw Minimum interval between applications: 7 days Minimum application volumes: 10 gallons/Acre (Ground); 2 gallons/Acre (Aerial) Maximum SIVANTO prime allowed per year: 28.0 fluid ounces/Acre (0.365 lb AI/Acre).	

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937)

Sivanto is a trademark of Bayer.

RESTRICTED USE PESTICIDE

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for those uses covered by the certified applicator's certification.

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee)
AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.



EPA Reg No: 279-3441

2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY IN
ARKANSAS, CONNECTICUT, DELAWARE, IDAHO, ILLINOIS,
INDIANA*, IOWA, KENTUCKY, MAINE, MARYLAND,
MASSACHUSETTS, MICHIGAN, MINNESOTA, MISSOURI, NEW
HAMPSHIRE, NEW JERSEY, NEW YORK, OHIO, OREGON,
RHODE ISLAND, VERMONT, VIRGINIA, WEST VIRGINIA,
WASHINGTON, WISCONSIN.

THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL
DIRECTIONS FOR THE USE WHICH ARE RECOMMENDED BY
FMC CORPORATION UNDER FIFRA SECTION 2(ee).

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT
IN A MANNER INCONSISTENT WITH ITS LABELING.

CROP	PEST	RATE OF APPLICATION
Apples	Mites European Red Mite Two-Spotted Spider Mite	19 fl oz/A (0.037 lbs ai/A)

DIRECTIONS FOR USE:

Broadcast ground only (minimum 40 gallons per acre). Begin applications at pink and/or petal fall through first cover to protect developing fruit and foliage from the target insect pest and mite species common to the production area. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels. Avoid applications when honey bees are actively foraging by applying during the early morning or evening hours.

ADJUVANTS

Surfactants can improve pest control. Do not use binder or sticker type surfactants. Vegetable oil can be used as a spray additive (1 qt/A). Use of a horticulture spray oil fewer than 14 days before or after applying Captan® or after sulfurcontaining products can result in phytotoxicity and crop loss. A nonionic surfactant is recommended to avoid fruit injury such as russetting on certain varieties.

Spider Mites: For better residual control of spider mites, applications should be made as close as possible to petal fall on newer leaves. Applications for spider mite control should be limited to a period from petal fall through 6 weeks following petal fall. Heavy mite pressure may need additional miticide for effective control. Apply before mites reach a threshold of 5 per leaf.

RESTRICTIONS:

- Do not apply more than 19 fl oz/A (0.037 lb ai/A) per application
- Do not apply more than 38 fl oz/A (0.15 lb ai/A) annually
- Do not make applications less than 21 days apart.
- PHI: 28 day
- Do not make more than 2 consecutive applications of Gladiator Insecticide/Miticide or any other product containing avermectin or zeta-cypermethrin. (Refer to max tables on section 3 label when making additional applications of other products containing either zeta-cypermethrin or avermectin)
- Do not apply using aerial applications.
- **Grazing Restrictions:** Do not allow livestock to graze or feed treated foliage to livestock

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED GLADIATOR® INSECTICIDE/MITICIDE (EPA REG. NO. 279-3441) LABEL MUST BE FOLLOWED. THIS RECOMMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

*Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist.



FMC Corporation
Agriculture Products Group
1735 Market Street
Philadelphia, PA 19103

Gladiator Insecticide/Miticide is a Restricted Use Pesticides.
Gladiator and FMC are registered trademarks of FMC Corporation.
©2016 FMC Corporation

2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



DUPONT™ GLEAN® XP herbicide

EPA Reg. No. 352-653

DUPONT™ ALLY® XP herbicide

EPA Reg. No. 352-435

DUPONT™ GR1™ herbicide

EPA Reg. No. 352-884

FOR POSTEMERGENCE WEED CONTROL OF BROADLEAF AND GRASSY WEEDS IN WINTER WHEAT and TRITICALE

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

GLEAN® XP herbicide may be tank mixed at 0.33 ounces per acre with 0.084 ounces per acre of ALLY® XP herbicide and 2.0 ounces per acre of GR1™ herbicide for the control of broadleaf and grassy weeds when applied postemergence to winter wheat and triticale in the fall or spring to 3-leaf stage but prior to jointing, when the majority of weeds have emerged and are actively growing. Refer to both labels for an entire lists of weeds controlled.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or DuPont 2(ee) instructions.

Read and follow all applicable directions, restrictions and precautions on the EPA-registered label for each product.

IMPORTANT PRECAUTIONS:

Refer to the GLEAN® XP herbicide, ALLY® XP herbicide, GR1™ herbicide and any tank mix partner labels for information on weed species controlled, additional use directions, restrictions, rotational crop intervals and precautions. The most restrictive provision on any label shall apply.

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2015 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A

R-1655 021616-2 02-16-16 C156

Page 1 of 1

Expires 12/31/18

2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



TANK MIXES OF:

DUPONT™ EXPRESS® herbicide (with TotalSol® soluble granules)

EPA Reg. No. 352-632

DUPONT™ ALLY® XP Herbicide

EPA Reg. No. 352-435

PLUS

DUPONT™ DICAMBA XP Herbicide

EPA Reg. No. 7969-140-352

OR

DUPONT™ BL1 Herbicide

EPA Reg. No. 42750-271-352

FOR POSTEMERGENCE WEED CONTROL OF BROADLEAF WEEDS IN WHEAT, BARLEY, TRITICALE and FALLOW

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

EXPRESS® herbicide (with TotalSol® soluble granules) may be tank mixed at 0.37 ounces per acre with ALLY® XP herbicide at 0.03 ounces per acre plus 1.80 ounces per acre of Dicamba XP herbicide or DuPont™ BL1 herbicide for the control of broadleaf weeds when applied postemergence to winter wheat in the fall or spring after the 2-leaf stage but prior to the jointing stage, when the majority of weeds have emerged and are actively growing. Spring applications to spring wheat must be made after the 2-leaf stage before the wheat exceeds the 6-leaf stage, and spring barley exceeds the 4-leaf stage, when the majority of weeds have emerged and are actively growing. Fall applications to triticale should be applied after the 2-leaf stage but prior to jointing, while spring seeded triticale applications must be made after the 2-leaf stage before the triticale reaches the 6-leaf stage. Application of this combination should not be made to small grains underseeded to legumes as severe legume injury or death will occur. Fallow applications should be made in the spring, summer, or fall when the majority of weeds have emerged and are actively growing. Refer to all labels for an entire lists of weeds controlled.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or DuPont 2(ee) instructions.

Read and follow all applicable directions, restrictions and precautions on the EPA-registered label for each product.

IMPORTANT PRECAUTIONS:

Refer to the EXPRESS® herbicide (with TotalSol® soluble granules), ALLY® XP herbicide, Dicamba XP herbicide, DuPont™ BL1 herbicide and any tank mix partner labels for information on weed species controlled, additional use directions, restrictions, rotational crop intervals and precautions. The most restrictive provision on any label shall apply.

© 2015 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2015 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A

2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



TANK MIXES OF:

DUPONT™ HARMONY® SG Herbicide
(with TotalSol® soluble granules)

EPA Reg. No. 352-633

DUPONT™ EXPRESS® Herbicide (with TotalSol® soluble granules)

EPA Reg. No. 352-632

PLUS

DUPONT™ DICAMBA XP Herbicide

EPA Reg. No. 7969-140-352

OR

DUPONT™ BL1 Herbicide

EPA Reg. No. 42750-271-352

FOR POSTEMERGENCE WEED CONTROL OF BROADLEAF WEEDS IN WHEAT, WINTER BARLEY, TRITICALE, and FALLOW

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

HARMONY® SG herbicide (with TotalSol® soluble granules) may be tank mixed at 0.51 ounces per acre with 0.39 ounces per acre of EXPRESS® herbicide (with TotalSol® soluble granules) plus 2.88 ounces per acre of Dicamba XP herbicide or DuPont™ BL1 herbicide for the control of broadleaf weeds when applied postemergence to winter wheat, winter barley, and triticale. Applications should be made in the fall or spring to winter wheat after the 2-leaf stage but prior to the jointing stage. Applications to winter barley must be made prior to the jointing stage. Spring applications to spring wheat must be made after the 2-leaf stage but before the wheat exceeds the 6-leaf stage. Fall applications to triticale should be applied after the 2-leaf stage but prior to jointing, while spring seeded triticale applications must be made after the 2-leaf stage but before the triticale reaches the 6-leaf stage. All applications should be made when the majority of weeds have emerged and are actively growing. Application of this combination should not be made to small grains underseeded to legumes as severe legume injury or death will occur. Fallow applications should be made in the spring, summer, or fall when the majority of weeds have emerged and are actively growing. Refer to all labels for an entire lists of weeds controlled.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or DuPont 2(ee) instructions.

Read and follow all applicable directions, restrictions and precautions on the EPA-registered label for each product.

© 2015 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A

IMPORTANT PRECAUTIONS:

Refer to the HARMONY® SG herbicide (with TotalSol® soluble granules), EXPRESS® herbicide (with TotalSol® soluble granules), Dicamba XP herbicide, DuPont™ BLI herbicide and any tank mix partner labels for information on weed species controlled, additional use directions, restrictions, rotational crop intervals and precautions. The most restrictive provision on any label shall apply.

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2015 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A

2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



TANK MIXES OF:

DUPONT™ EXPRESS® herbicide (with TotalSol® soluble granules)

EPA Reg. No. 352-632

DUPONT™ ALLY® XP Herbicide

EPA Reg. No. 352-435

PLUS

DUPONT™ DICAMBA XP Herbicide

EPA Reg. No. 7969-140-352

OR

DUPONT™ BL1 Herbicide

EPA Reg. No. 42750-271-352

FOR POSTEMERGENCE WEED CONTROL OF BROADLEAF WEEDS IN WHEAT, WINTER BARLEY, TRITICALE and FALLOW

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

EXPRESS® herbicide (with TotalSol® soluble granules) may be tank mixed at 0.37 ounces per acre with ALLY® XP herbicide at 0.03 ounces per acre plus 2.88 ounces per acre of Dicamba XP herbicide or DuPont™ BL1 herbicide for the control of broadleaf weeds when applied postemergence to winter wheat and winter barley in the fall or spring after the 2-leaf stage but prior to the jointing stage when the majority of weeds have emerged and are actively growing. Spring applications to spring wheat must be made after the 2-leaf stage before the wheat exceeds the 6-leaf stage, when the majority of weeds have emerged and are actively growing. Fall applications to triticale should be applied after the 2-leaf stage but prior to jointing, while spring seeded triticale applications must be made after the 2-leaf stage before the triticale reaches the 6-leaf stage. Application of this combination should not be made to small grains underseeded to legumes as severe legume injury or death will occur. Fallow applications should be made in the spring, summer, or fall when the majority of weeds have emerged and are actively growing. Refer to all labels for an entire lists of weeds controlled.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or DuPont 2(ee) instructions.

Read and follow all applicable directions, restrictions and precautions on the EPA-registered label for each product.

IMPORTANT PRECAUTIONS:

Refer to the EXPRESS® herbicide (with TotalSol® soluble granules), ALLY® XP herbicide, Dicamba XP herbicide, DuPont™ BL1 herbicide and any tank mix partner labels for information on weed species controlled, additional use directions, restrictions, rotational crop intervals and precautions. The most restrictive provision on any label shall apply.

© 2015 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2015 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A

2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



TANK MIXES OF:

DUPONT™ HARMONY® SG herbicide (with TotalSol® soluble granules)

EPA Reg. No. 352-633

DUPONT™ EXPRESS® herbicide (with TotalSol® soluble granules)

EPA Reg. No. 352-632

DUPONT™ ALLY® XP Herbicide

EPA Reg. No. 352-435

PLUS

DUPONT™ DICAMBA XP Herbicide

EPA Reg. No. 7969-140-352

OR

DUPONT™ BL1 Herbicide

EPA Reg. No. 42750-271-352

FOR POSTEMERGENCE WEED CONTROL OF BROADLEAF WEEDS IN WHEAT, BARLEY, TRITICALE and FALLOW

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

HARMONY® SG herbicide (with TotalSol® soluble granules) may be tank mixed at 0.272 ounces per acre with EXPRESS® herbicide (with TotalSol® soluble granules) at 0.137 ounces per acre plus ALLY® XP herbicide at 0.091 ounces per acre plus 1.80 ounces per acre of Dicamba XP herbicide or DuPont™ BL1 herbicide for the control of broadleaf weeds when applied postemergence to winter wheat in the fall or spring after the 2-leaf stage but prior to the jointing stage, when the majority of weeds have emerged and are actively growing. Spring applications to spring wheat must be made after the 2-leaf stage before the wheat exceeds the 6-leaf stage, and spring barley exceeds the 4-leaf stage, when the majority of weeds have emerged and are actively growing. Fall applications to triticale should be applied after the 2-leaf stage but prior to jointing, while spring seeded triticale applications must be made after the 2-leaf stage before the triticale reaches the 6-leaf stage. Application of this combination should not be made to small grains underseeded to legumes as severe legume injury or death will occur. Fallow applications should be made in the spring, summer, or fall when the majority of weeds have emerged and are actively growing. Refer to all labels for an entire lists of weeds controlled.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or DuPont 2(ee) instructions.

© 2015 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A

Read and follow all applicable directions, restrictions and precautions on the EPA-registered label for each product.

IMPORTANT PRECAUTIONS:

Refer to the HARMONY® SG herbicide (with TotalSol® soluble granules), EXPRESS® herbicide (with TotalSol® soluble granules), ALLY® XP herbicide, Dicamba XP herbicide, DuPont™ BL1 herbicide and any tank mix partner labels for information on weed species controlled, additional use directions, restrictions, rotational crop intervals and precautions. The most restrictive provision on any label shall apply.

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2015 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A