

2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



DUPONT™ EXPRESS® XP HERBICIDE

EPA Reg. No. 352-509

DUPONT™ HARMONY® GT XP HERBICIDE

EPA Reg. No. 352-446

FOR POSTEMERGENCE WEED CONTROL OF BROADLEAF WEEDS IN WHEAT (including durum), BARLEY, OATS, TRITICALE, and FALLOW and BURNDOWN

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

EXPRESS® XP herbicide may be tank mixed at 0.135 ounces per acre to 0.33 ounces per acre with HARMONY® GT XP herbicide at 0.135 ounces per acre to 0.335 ounces per acre, in a 1 to 1 ratio, for control of broadleaf weeds when applied postemergence to wheat (including durum), barley, and triticale in the 2-leaf stage but before flag leaf is visible and fallow or burndown in the spring, summer or fall when the majority of weeds have emerged and are actively growing. EXPRESS® XP herbicide may be tank mixed at 0.133 ounces per acre with HARMONY® GT XP herbicide at 0.135 ounces per acre, for control of broadleaf weeds when applied postemergence to winter oats in the 2-leaf stage but before flag leaf is visible or spring oats in the 3-leaf stage but before jointing. Refer to both labels for an entire lists of weeds controlled.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or DuPont 2(ee) instructions.

Read and follow all applicable directions, restrictions and precautions on the EPA-registered label for each product.

IMPORTANT PRECAUTIONS:

Refer to the EXPRESS® XP herbicide, HARMONY® GT XP herbicide, and any tank mix partner labels for information on weed species controlled, additional use directions, restrictions, rotational crop intervals and precautions. The most restrictive provision on any label shall apply.

(Replaces: R-1644 092315-2 09-29-15 C-152)

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2016 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Road, Wilmington, DE 19805 U.S.A.
Unless indicated; trademarks with ®; TM or SM are trademarks of DuPont or its affiliates. © 2015 DuPont



Section 2(ee) Recommendation

Date: April 22, 2016

Product: Acuron® Herbicide
EPA Reg. No. 100-1466

Use: Impregnation or coating onto dry bulk fertilizer

State: IA, IL, MN, ND, SD, WI

Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- The user must refer to the federally approved labels for the above noted product and read and follow all directions for use, restrictions, and precautions.
- The user should have this recommendation in its possession at the time of use.
- This recommendation for use of this product is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.

Acuron Herbicide may be impregnated or coated onto dry bulk fertilizers including ammonium phosphate-sulfate, ammonium sulfate (AMS), diammonium phosphate (DAP), monoammonium phosphate (MAP), potash (potassium chloride), potassium sulfate, urea, or blends of these dry bulk fertilizer types.

When applying Acuron Herbicide on dry bulk fertilizer, follow all directions for use and precautions on the product label regarding target crops, application rate, timing of application and all precautions and restrictions.

All individual state regulations relating to dry bulk granular fertilizer blending, registration, labeling, and application are the responsibility of the mixer and applicator.

Apply a minimum of 200 lb of dry bulk fertilizer impregnated with Acuron Herbicide at the recommended broadcast rate per acre. For best results, apply the mixture uniformly to the soil with properly calibrated equipment immediately after blending. Uniform application of the blended fertilizer/herbicide mixture is essential to prevent possible crop injury and achieve weed control. Non-uniform application will result in unsatisfactory weed control. In areas where tillage is practiced, a shallow incorporation of the blended fertilizer/herbicide mixture is recommended for improved weed control.

Calculate amount of Acuron Herbicide needed by the following formula:

$$\frac{2,000}{\text{lb of fertilizers per acre}} \times \frac{\text{qt/A of Acuron Herbicide}}{\text{qt of Acuron Herbicide per ton of fertilizer}} =$$

Blended Mixing Instructions

Prepare the fertilizer/herbicide mixture by using any closed drum, belt, ribbon, or other commonly used dry bulk fertilizer blender. Nozzles used to spray Acuron Herbicide onto the fertilizer must be placed to provide uniform spray coverage. Care should be taken to aim the spray directly onto the fertilizer and avoid spraying the walls of the blender.

If the fertilizer/herbicide blend is too wet for uniform application, adding a drying agent is recommended. Add the drying agent slowly to the fertilizer/herbicide blend until the mixture is suitable for uniform application. The amount of drying agent needed will depend on fertilizer type, Acuron Herbicide application rate and amount of fertilizer used.

Apply the fertilizer/herbicide blend immediately following impregnation.

Pneumatic (Compressed Air) Application:

Acuron Herbicide may be applied through pneumatic applicators, whether the fertilizer/herbicide mixture is blender-mixed or on-board fertilizer impregnation system.

Acuron Herbicide must not be mixed with any other liquid or dry material in on-board fertilizer impregnation system tanks. Use high quality fertilizer with a minimum of fines when applying Acuron Herbicide with on-board impregnation equipment.

Drying agents are not recommended for use with on-board impregnation systems.

Precautions

- (1) To avoid potential for explosion, do not impregnate Acuron Herbicide onto ammonium nitrate, potassium nitrate, or sodium nitrate either alone or in blends with other fertilizers.
- (2) Do not impregnate Acuron Herbicide onto single super phosphate or triple superphosphate fertilizers.
- (3) Do not impregnate Acuron Herbicide on straight unadulterated agricultural limestone, since absorption will not be achieved.

Acuron® and the Syngenta logo are trademarks of a Syngenta Group Company.

©2016 Syngenta

2(ee) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Expiration Date: April 12, 2021

Label Code: IA1466040CC0416



Bayer CropScience, LP
 P.O. Box 12014
 2 T.W. Alexander Drive
 Research Triangle Park, North Carolina 27709
 1-866-99BAYER (1-866-992-2937)

STREAMLINE HERBICIDE®

FOR CONTROL OR SUPPRESSION OF
 ADDITIONAL WEEDS AND BRUSH IN NON-
 CROP AREAS

EPA Reg. No. 432-1570

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN ALL STATES EXCEPT NEW HAMPSHIRE, NEW YORK, AND TEXAS

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

PRODUCT INFORMATION

Streamline Herbicide is recommended for the control of listed weeds and brush grown in non-cropland areas as described in the product information section of the Streamline Herbicide federal label.

WEEDS CONTROLLED:

4.75 to 9.5 ounces per acre

Hakweed, orange	<i>Hieracium aurantiacum</i>
Locust, honey	<i>Gleditsia triacanthos</i>
Snakeweed, broom	<i>Gutierrezia sarothrae</i>
Toadflax, dalmation	<i>Linaria dalmatica</i>

7.5 to 11.5 ounces per acre

Broom, scotch	<i>Cytisus scoparius</i>
---------------	--------------------------

BRUSH CONTROLLED:

BRUSH		Rate High Volume Foliar Ounces/100 gallon	Rate Broadcast Foliar Ounces/acre
Pine, loblolly	<i>Pinus taeda</i>	9.5-11.5	11.5
Pine, Virginia ^{1,2}	<i>Pinus virginia</i>	9.5-11.5	11.5
Privet, Chinese	<i>Ligustrum sinense</i>	9.5-11.5	11.5
Rabbitbrush, rubber	<i>Ericameria nauseosa</i>	9.5-11.5	11.5
Sweetgum ^{1,2}	<i>Liquidambar styraciflua</i>	9.5-11.5	11.5

- 1- Suppression: a visual reduction in weed competition (reduced population or vigor) as compared to an untreated area
- 2- See specific weed section



Bayer CropScience, LP
P.O Box 12014
2 T.W Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

STREAMLINE HERBICIDE®

FOR CONTROL OR SUPPRESSION OF
ADDITIONAL WEEDS AND BRUSH IN NON-
CROP AREAS

EPA Reg. No. 432-1570

FIFRA 2(ee) Recommendation

SPECIFIC WEED DIRECTIONS:

Virginia Pine: For complete control, tankmix "Krenite" at 1.5 gallons per acre or glyphosate at 5.4 pounds active ingredient per acre (1 gallon of a 5.4 pound per gallon material of 1.33 gallons of a 4 pound gallon material).

Sweetgum: For complete control, tankmix "Krenite" at 1.5 gallons per acre or glyphosate at 5.4 pounds active ingredient per acre (1 gallon of a 5.4 pound gallon material or 1.333 gallons of a 4 pound gallon material).

Use a suitable adjuvant with Streamline Herbicide as listed in the adjuvant section of the Streamline federal label. Do not apply more than 11.5 ounces per broadcast per year.

Use the higher spray volumes, herbicide and adjuvant rate for heavy weed infestations and hard to control species.

Through spray coverage of all brush foliage is necessary for best control.

Before using Streamline Herbicide, refer to the product label for additional information and precautions. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on EPA registered label.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

*As with any herbicide product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*



Bayer CropScience, LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

Perspective Herbicide®

For the control of broadleaf weeds in non-crop sites.

EPA Reg. No. 432-1569

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN ALL STATES EXCLUDING NEW HAMPSHIRE, NEW YORK, AND TEXAS

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label. Always read this FIFRA 2 (ee) bulletin and the entire label on the product container before using this product. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product in any way that will contact workers of other persons, either directly or through drift. Only handlers may be in the area during application.

Bayer will not be responsible for losses or damages resulting from the use Perspective Herbicide in any manner not specifically instructed by Bayer. User assumes all risks associated with non-labeled uses.

PRODUCT INFORMATION

Perspective Herbicide is a dispersible granule that is mixed in water and applied as a spray. Perspective Herbicide may be applied by ground equipment only, except for the rights-of-ways which may also be applied by helicopter, for control of broadleaf weeds, including many terrestrial and riparian invasive and noxious weeds. Perspective is registered for general weed control on private, public and military lands as follows: uncultivated non-agricultural areas (such as airports, highway, railroad, and utility rights-of-way, sewage disposals areas, etc.); uncultivated agricultural areas- non-crop producing (such as farmyards, fuel storage areas, fence rows, non-irrigation ditchbanks, barrier strips, etc.); industrial sites- outdoor (such as lumberyards, pipeline and tank farms, etc.); and natural areas (such as wildlife management areas, wildlife openings and wildlife habitats). Perspective Herbicide may be used for the release of restoration of native perennial grasses and in established, industrial turf grasses.

This product may be applied to terrestrial non-crop sites and unimproved turf sites that contain areas of temporary surface water caused by collection of water, in equipment ruts, or in other depressions created by management activities. It is permissible to treat intermittently flooded low lying sites, seasonally dry flood plains and transitional areas between upland and lowland sites when no water is present. It is also permissible to treat marshes, swamps and bogs after water has receded, as well as seasonally dry flood deltas. Perspective Herbicide may be applied up to the waters edge. Do not apply directly to water.

Perspective Herbicide provides preemergence and/or postemergence control of the broadleaf weeds species listed in the weeds controlled section of the label. For perennial species on the label, a postemergence application must be used. For best postemergence performance, add a high quality adjuvant to the spray solution. Excessive wetting of the target plant is not necessary but good spray coverage of the target plant is needed for best results. Do not apply more than 11 ounces of product per acre per year. Do not apply more than three times per year.



Bayer CropScience, LP
P.O Box 12014
2 T.W Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

Perspective Herbicide®

For the control of broadleaf weeds in non-crop sites.

EPA Reg. No. 432-1569

FIFRA 2(ee) Recommendation

WEEDS CONTROLLED

RATE

Hawkweed, orange

RATE

Locust, honey

Thistle, Russian

Toadflax, dalmation

Toadflax, yellow

3.0 to 4.5 ounces per acre

Hieracium aurantiacum

4.75 to 8 ounces per acre

Gleditsia triacanth

Salsola kali

Linaria dalmatica

Linaria vulgaris

IMPORTANT

Refer to Perspective Herbicide and any tank-mix partner labels for information on adjuvants; weed species controlled; additional use directions; rotations crop intervals and precautions. The most restrictive provision on any label should apply.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

*As with any herbicide product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*



Section 2(ee) Recommendation

Date: April 14, 2016
Product: Acuron® Flexi
EPA Reg. No. 100-1568
Use: Impregnation or coating onto dry bulk fertilizer
State: IA, IL, MN, ND, SD, WI

Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- The user must refer to the federally approved labels for the above noted product and read and follow all directions for use, restrictions, and precautions.
- The user should have this recommendation in its possession at the time of use.
- This recommendation for use of this product is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.

Acuron Flexi may be impregnated or coated onto dry bulk fertilizers including ammonium phosphate-sulfate, ammonium sulfate (AMS), diammonium phosphate (DAP), monoammonium phosphate (MAP), potash (potassium chloride), potassium sulfate, urea, or blends of these dry bulk fertilizer types.

When applying Acuron Flexi on dry bulk fertilizer, follow all directions for use and precautions on the product label regarding target crops, application rate, timing of application and all precautions and restrictions.

All individual state regulations relating to dry bulk granular fertilizer blending, registration, labeling, and application are the responsibility of the mixer and applicator.

Apply a minimum of 200 lb of dry bulk fertilizer impregnated with Acuron Flexi at the recommended broadcast rate per acre. For best results, apply the mixture uniformly to the soil with properly calibrated equipment immediately after blending. Uniform application of the blended fertilizer/herbicide mixture is essential to prevent possible crop injury and achieve weed control. Non-uniform application will result in unsatisfactory weed control. In areas where tillage is practiced, a shallow incorporation of the blended fertilizer/herbicide mixture is recommended for improved weed control.

Calculate amount of Acuron Flexi needed by the following formula:

$$\frac{2,000}{\text{lb of fertilizers per acre}} \times \text{qt/A of Acuron Flexi} = \text{qt of Acuron flexi per ton of fertilizer}$$

Blended Mixing Instructions

Prepare the fertilizer/herbicide mixture by using any closed drum, belt, ribbon, or other commonly used dry bulk fertilizer blender. Nozzles used to spray Acuron Flexi onto the fertilizer must be placed to provide uniform spray coverage. Care should be taken to aim the spray directly onto the fertilizer and avoid spraying the walls of the blender.

If the fertilizer/herbicide blend is too wet for uniform application, adding a drying agent is recommended. Add the drying agent slowly to the fertilizer/herbicide blend until the mixture is suitable for uniform application. The amount of drying agent needed will depend on fertilizer type, Acuron Flexi application rate and amount of fertilizer used.

Apply the fertilizer/herbicide blend immediately following impregnation.

Pneumatic (Compressed Air) Application:

Acuron Flexi may be applied through pneumatic applicators, whether the fertilizer/herbicide mixture is blender-mixed or on-board fertilizer impregnation system.

Acuron Flexi must not be mixed with any other liquid or dry material in on-board fertilizer impregnation system tanks. Use high quality fertilizer with a minimum of fines when applying Acuron Flexi with on-board impregnation equipment.

Drying agents are not recommended for use with on-board impregnation systems.

Precautions

- (1) To avoid potential for explosion, do not impregnate Acuron flexi onto ammonium nitrate, potassium nitrate, or sodium nitrate either alone or in blends with other fertilizers.
- (2) Do not impregnate Acuron Flexi onto single super phosphate or triple superphosphate fertilizers.
- (3) Do not impregnate Acuron Flexi on straight unadulterated agricultural limestone, since absorption will not be achieved.

Acuron® and the Syngenta logo are trademarks of a Syngenta Group Company.

©2016 Syngenta

2(ee) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Expiration Date: April 14, 2021

Label Code: IA1568039AC0416



Bayer CropScience

Bayer CropScience LP
 P.O. Box 12014
 2 T.W. Alexander Drive
 Research Triangle Park, North Carolina 27709
 1-866-99BAYER (1-866-992-2937)

SIVANTO™ prime
 EPA Reg. No. 264-1141

Reduced rates for foliar applications in Pearl Millet and Sorghum for control of Sugarcane Aphid.

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF ILLINOIS

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. This product bulletin should be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PEARL MILLET; SORGHUM – FOLIAR

Pest Controlled	Product Rate (fl oz/A)
Sugarcane Aphid	4.0 – 7.0
Foliar Application Restrictions: Pre-Harvest Interval (PHI): 7 days - forage; 21 days - dried grain, stover or straw Minimum interval between applications: 7 days Minimum application volumes: 10 gallons/Acre (Ground); 2 gallons/Acre (Aerial) Maximum SIVANTO prime allowed per year: 28.0 fluid ounces/Acre (0.365 lb AI/Acre).	

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937)

Sivanto is a trademark of Bayer.

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.



EPA REG No. 84876-1-279

2(ee) Recommendation
FOR DISTRIBUTION AND USE ONLY IN CONNECTICUT, DELAWARE, IDAHO, ILLINOIS, INDIANA*, KENTUCKY, MAINE, MARYLAND, MICHIGAN, NEW HAMPSHIRE, NEW JERSEY, NEW YORK, OHIO, OREGON, PENNSYLVANIA, RHODE ISLAND, VERMONT, VIRGINIA, WEST VIRGINIA, WASHINGTON, WISCONSIN
THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL DIRECTIONS FOR USE WHICH ARE RECOMMENDED BY FMC CORPORATION UNDER FIFRA SECTION 2(ee)
IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH IT LABELING

CROP	PEST	RATE OF APPLICATION
Grape	[Suppression] Sour Rot (<i>Acetobacter spp.</i>)	24.4 oz – 36.6 oz/A (0.40 – 0.60 lbs ai /A)
DIRECTIONS FOR USE: Sour Rot [Suppression]: Begin applications at veraison and / or when sugar concentration of grape berries has reached 8 to 14%.		
PRECAUTIONS: Alternate applications with another effective fungicide with a different mode of action. Fracture fungicides requires 2 – 4 hours drying time on plant foliage for the active ingredient to fix into the plant tissue before rain or irrigation occurs. If during the next 12 hours it rains significantly, a new application will be needed during the next 4 days.		
RESTRICTIONS: <ul style="list-style-type: none"> • Do not make more than 5 foliar applications of Fracture fungicide per crop cycle • Do not make more than two sequential applications of Fracture Fungicide before alternating to a labeled fungicide with a different mode of action. • Do not harvest until 1 day after last application • Follow most restrictive plant back intervals, crop rotations, and other intervals when tank mixing with other products 		
ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED FRACTURE FUNGICIDE LABEL (EPA REG. NO. 84876-1-279) MUST BE FOLLOWED. THIS RECOMMENDATION SHOULD BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.		

* Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist.

FMC
 FMC Corporation
 Agriculture Products Group
 1735 Market Street
 Philadelphia, PA 19103

Fracture Fungicide - FMC Trademark
 ©2016 FMC Corporation
 Version: 2016-3-18



Section 2(ee) Recommendation

Date: March 22, 2016

Product: Segovis®, EPA Reg. No. 100-1533

Use: Addition of Downy Mildew as a Pathogen Controlled by Drench Application to Container Grown Ornamental Plants

State: AK, AL, AR, AZ, CO, CT, DE, FL, GA, IA, ID, IL, IN, KS, KY, LA, MA, MD, ME, MI, MN, MO, MS, MT, NC, ND, NE, NH, NJ, NM, NV, OH, OK, PA, RI, SC, SD, TN, UT, VA, VT, WA, WI, WV, WY

Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- The user must refer to the federally approved labels for the above noted product and read and follow all directions for use, restrictions, and precautions.
- The user should have this recommendation in its possession at the time of use.
- Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist.
- This recommendation for use of this product is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.

Crop	Disease/ Pathogen	Rate fl oz/100 gal	Application Interval/ Timing	Instructions
Breeding crops Bulb crops (including Calla Lilies, Easter Lilies, Gladiolas and Caladiums) Cut flowers Evergreens (including conifers) Flowering plants Flowers grown for seed production Foliage plants Ground covers Non-bearing fruit trees Non-bearing nut trees Non-bearing vines Ornamental trees and shrubs Perennial plants Pot and bedding plants	Downy Mildew	1.0-3.2	5-14 days	Apply as a drench at volumes specified on the Segovis Section 3 product label.

Segovis® and the Syngenta logo are trademarks of a Syngenta Group Company.
©2016 Syngenta

2(ee) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Expiration Date: March 22, 2021

Label Code: AK1533004AC0316

